

Original Article

Timing and route of contamination of hospitalized patient rooms with healthcare-associated pathogens

Sarah N. Redmond BS¹, Basya S. Pearlmutter BS², Yilen K. Ng-Wong MD², Heba Alhmidi MD², Jennifer L. Cadnum BS², Sandra Y. Silva MD³, Brigid M. Wilson PhD⁴ and Curtis J. Donskey MD^{1,4}

¹Case Western Reserve University School of Medicine, Cleveland, Ohio, ²Research Service, Louis Stokes Cleveland Veterans' Affairs Medical Center, Cleveland, Ohio, ³Clinical and Translational Science Program, School of Medicine, Case Western Reserve University and ⁴Geriatric Research, Education, and Clinical Center, Louis Stokes Cleveland VA Medical Center, Cleveland, Ohio

Abstract

Objective: To investigate the timing and routes of contamination of the rooms of patients newly admitted to the hospital.

Design: Observational cohort study and simulations of pathogen transfer.

Setting: A Veterans' Affairs hospital.

Participants: Patients newly admitted to the hospital with no known carriage of healthcare-associated pathogens.

Methods: Interactions between the participants and personnel or portable equipment were observed, and cultures of high-touch surfaces, floors, bedding, and patients' socks and skin were collected for up to 4 days. Cultures were processed for *Clostridioides difficile*, methicillin-resistant *Staphylococcus aureus* (MRSA), and vancomycin-resistant enterococci (VRE). Simulations were conducted with bacteriophage MS2 to assess plausibility of transfer from contaminated floors to high-touch surfaces and to assess the effectiveness of wearing slippers in reducing transfer.

Results: Environmental cultures became positive for at least 1 pathogen in 10 (59%) of the 17 rooms, with cultures positive for MRSA, *C. difficile*, and VRE in the rooms of 10 (59%), 2 (12%), and 2 (12%) participants, respectively. For all 14 instances of pathogen detection, the initial site of recovery was the floor followed in a subset of patients by detection on sock bottoms, bedding, and high-touch surfaces. In simulations, wearing slippers over hospital socks dramatically reduced transfer of bacteriophage MS2 from the floor to hands and to high-touch surfaces.

Conclusions: Floors may be an underappreciated source of pathogen dissemination in healthcare facilities. Simple interventions such as having patients wear slippers could potentially reduce the risk for transfer of pathogens from floors to hands and high-touch surfaces.

(Received 13 July 2020; accepted 1 December 2020; electronically published 12 January 2021)

Patients admitted to healthcare facilities are at risk to acquire colonization or infection with healthcare-associated pathogens such as *Clostridioides difficile*, methicillin-resistant *Staphylococcus aureus* (MRSA), and vancomycin-resistant enterococci (VRE). The hands of healthcare personnel are generally considered the most important source of pathogen transmission.¹ Contaminated surfaces and equipment have also been implicated in transmission, either through direct contact with patients or indirectly when hands of personnel transfer pathogens from surfaces or equipment to patients.^{1–4} Admission to a room previously occupied by a colonized or infected patient increases the risk for subsequent room occupants to acquire the same organism, suggesting acquisition from inadequately cleaned surfaces.^{1,2} Contaminated shared electronic thermometers and temperature probes have been linked to transmission of *C. difficile*, VRE, and the emerging fungal pathogen *Candida auris*.^{6–8} Finally,

some recent studies have suggested that floors might be an underappreciated source of pathogen transmission.^{9–13} For example, a nonpathogenic virus inoculated onto floors in patient rooms disseminated rapidly to the hands of patients and to surfaces in the room.⁹

To develop effective control measures, a better understanding of sources of pathogen exposure in healthcare facilities is needed. In the current study, we investigated the timing and routes of contamination of the rooms of patients newly admitted to the hospital, focusing on individuals with negative nares screen for MRSA and no known carriage of healthcare-associated pathogens. We hypothesized that direct interactions with healthcare personnel or medical equipment would be associated with recovery of pathogens from surfaces and patients.

Methods

Study setting

The Cleveland Veterans' Affairs (VA) Medical Center is a 215-bed acute-care hospital. All admitted patients are screened for nasal

Author for correspondence: Curtis J. Donskey, E-mail: Curtis.Donskey@va.gov

Cite this article: Redmond SN, et al. (2021). Timing and route of contamination of hospitalized patient rooms with healthcare-associated pathogens. *Infection Control & Hospital Epidemiology*, 42: 1076–1081, <https://doi.org/10.1017/ice.2020.1367>

carriage of MRSA. At the time of the study, hand hygiene compliance in the facility was measured at >85% based on observations by dedicated hand hygiene monitors.

Observational cohort study

We conducted an observational cohort study of patients newly admitted to single-patient rooms on medical-surgical wards. Patients with prior colonization or infection with MRSA, VRE, or *C. difficile* within the past year were excluded. The study was approved by the facility's institutional review board. Prior to enrollment, patient rooms received routine postdischarge cleaning by environmental services personnel followed by additional cleaning and disinfection by research personnel. Research personnel applied a sporicidal cleaner-disinfectant product containing 2,500 parts per million sodium hypochlorite on all high-touch surfaces and on the floor and allowed the surfaces to air dry. A low-pressure mercury ultraviolet-C (UV-C) light room decontamination device (UVDI-360 Room Sanitizer, Ultraviolet Devices, Santa Clarita, CA) was then run for two 5-minute cycles in the patient room (one on each side of the bed) and for one 5-minute cycle in the bathroom.¹⁴ Cultures of high-touch surfaces (bed rail, bedside table, call button, patients' bedside chart) and floors (5×10-cm areas adjacent to the patients' bed) were collected to assess effectiveness of cleaning and disinfection. If any of these preadmission cultures returned positive, the patient's data were excluded from the assessment of routes of room contamination.

After admission, interactions between the participating patients and personnel and portable equipment were observed for ~6 hours per day by research personnel. Any entry of personnel or equipment into the room with or without contact with the patient and/or environment was considered an interaction. The research personnel were primarily stationed outside the rooms but entered the room if needed to observe care interactions; if interactions could not be directly observed (eg, curtains pulled), research personnel would ask providers to describe the procedures being performed and the types of contact. Hospital personnel were aware that a study related to environmental contamination was being conducted but were not informed of the study goals. Cultures of high-touch surfaces (bed rails, bedside tables, call buttons, patients' bedside charts) and floors (5×10-cm areas adjacent to the patients' beds) as well as the hands, chest, groin, and sock bottoms of patients and bedding (ie, sheets and blankets at the base of the bed where feet are placed) were collected as previously described at the time of admission, 4 and 12 hours after admission, and at least once daily on subsequent days for up to 4 days if the patient remained hospitalized. Additional sets of cultures were collected if observed interactions involved direct or indirect contact between personnel or equipment and patients or environmental surfaces; the sites of contact were sampled (eg, patient's skin or surfaces or equipment would be sampled if contacted by personnel or patients during care activities).

Microbiology and molecular typing

BBL CultureSwabs (Becton Dickinson) premoistened with Dey Engle neutralizer were used for sample collection. Cultures were processed for *C. difficile*, MRSA, and VRE as previously described.¹⁵ For selected patients with MRSA isolates recovered from multiple sites, *spa* typing was performed.¹⁶

Use of simulations to test the effectiveness of wearing slippers in reducing transfer from floors to high-touch surfaces

Based on results of the hospital-based study implicating floors as a route of pathogen transmission, we hypothesized that having patients wear slippers would be effective in reducing transfer of pathogens from floors to high-touch surfaces. To test this hypothesis, we used simulations with the nonpathogenic virus bacteriophage MS2 comparing transfer when socks were worn versus when slippers were worn over the socks. The simulations were conducted in a simulated patient room.

Bacteriophage MS2 was prepared as previously described.¹⁷ A 1-mL suspension containing 10^8 plaque-forming units (PFU) of MS2 was applied to a 10-cm × 10-cm area of the floor adjacent to the bed and allowed to air dry. Two types of simulations were conducted to assess whether wearing slippers would reduce transfer of MS2 from the floor to high-touch surfaces in a simulated patient room containing a patient bed and bedside table. In the first simulation, research personnel walked through the area of contamination while wearing hospital patient socks or socks plus slippers. They removed the slippers (if worn) and lay down on the bed, touched the bedding in contact with their socks, and then touched the bedside table. The second simulation was identical to the first except the participants removed their socks after sitting on the bed and then touched the bedside table. BBL CultureSwabs were used to sample the floor, sock bottoms, hands, bedding, and tray table. Bacteriophage MS2 was quantified as previously described.¹⁷ The simulations were repeated 3 times.

Data analysis

Analysis of variance was used to compare the log₁₀ PFU of bacteriophage MS2 to socks, bedding, hands, and bedside tables. Data were analyzed using R version 3.5.0 software (R Foundation for Statistical Computing, Vienna, Austria).

Results

In total, 25 patient rooms were cleaned and disinfected in anticipation of recruitment of new admissions for enrollment. Of the 25 patients considered for enrollment, 8 (32%) declined participation. Of the 17 study participants, 16 (94%) were ambulatory. The mean age of the participants was 66.9 (range, 24–82) and 15 (88.2%) were male. Based on observations, the participants interacted with an average of 2.4 personnel and 0.4 portable devices per hour. The average number of personnel and portable equipment interactions observed per patient were 15 (range, 3–32) and 3.5 (range, 0–11). Figure 1 shows the number and types of personnel and portable devices that entered the rooms. The average length of patient participation was 2.2 days (range, 1–4).

All room cultures collected after cleaning and disinfection and before admission were negative (174 total). In total, 1,208 environmental and skin swab samples were obtained for culture after patient admission, including 418 that were collected based on observations of interactions between patients and personnel and/or portable equipment and 790 that were sampled at scheduled intervals. One or more environmental cultures became positive for at least 1 of the pathogens in 10 of the 17 (59%) rooms. MRSA, *C. difficile*, and/or VRE were recovered in rooms of 10 (59%), 2 (12%), and 2 (12%) participants, respectively. Table 1 shows the pathogens recovered from the different sampling sites for the 10 participants who had 1 or more positive room cultures.




Fig. 1. Frequency of interactions between 17 study participants and personnel (A) and shared portable equipment and fomites (B). The equipment and fomites were taken inside the room and either directly contacted patients or were contacted by personnel who subsequently touched the patients. Other personnel included social work, nutrition, pharmacy, dialysis, patient transport, and phlebotomy. Workstations were computerized workstations on wheels. Other equipment and fomites included portable vital signs equipment, electrocardiogram machine, echocardiogram machine, and transfer devices.

For all 14 instances where pathogens were recovered, the initial site of recovery was the floor.

Figure 2 shows the cumulative percentage of MRSA contamination in rooms and the percentage of patients remaining the hospital. For all 10 patients with positive MRSA cultures, contamination was initially detected on the floor after personnel entered the room; for 6 of the 10 (60%) patients, MRSA was detected on the floor within 8 hours of admission. No cultures of surfaces or patients' skin became positive for MRSA when samples were collected immediately after personnel and/or equipment were observed interacting with the subjects ($N = 418$ sets of cultures). MRSA was recovered from high-touch surfaces in 3 (17.6%) of the 17 rooms. In 2 (66.7%) of 3 instances when MRSA was recovered from high-touch surfaces, an isolate with the same spa type was recovered from a previous floor sample. In 1 instance when MRSA was recovered from high-touch surfaces, an MRSA isolate with a different ribotype was recovered from a previous floor sample. For 3 patients, MRSA isolates

recovered from the floor had the same spa type as isolates subsequently recovered from other sites (socks, bedding, and/or high touch surfaces).

Figure 3 shows the average concentration of bacteriophage MS2 recovered from different sites with the 2 simulation protocols. For both protocols, MS2 was transferred from the floor to all the subsequent sampling sites when socks were worn. When slippers were worn over the socks, contamination was significantly reduced on socks, bedding, hands, and bedside tables ($P < .01$). No MS2 was detected on the hands or bedside table when slippers were worn.

Discussion

The hands of personnel and contaminated surfaces and equipment are generally recognized as important vectors for transmission of healthcare-associated pathogens.¹⁻³ Infection control measures therefore emphasize efforts to improve hand hygiene and cleaning of surfaces and equipment. In the current study, we did not find

Table 1. Culture Results for the 10 Newly Admitted Patients with 1 or More Positive Cultures for Methicillin-Resistant *Staphylococcus aureus* (MRSA), *Clostridioides difficile*, and Vancomycin-Resistant Enterococci (VRE)

Patient No.	Length of Stay, h	Floor ^a	Socks	Bedding ^b	High-Touch Surfaces ^c	Skin ^d
1	27	MRSA	ND	ND
2	42	MRSA
3	26	MRSA	MRSA
4	49	MRSA
5	72	MRSA	MRSA	...	MRSA	...
6	96	MRSA
7	48	MRSA, <i>C difficile</i>	...	<i>C difficile</i>	MRSA	...
8	72	MRSA, VRE	...	VRE
9	96	MRSA, VRE, <i>C difficile</i>	MRSA	MRSA
10	45	MRSA


Note. ND, not done.

^aFloor cultures included 5x10-cm areas adjacent to the patients' beds.

^bBedding included sheets and blankets at the base of the bed where feet are placed.

^cHigh-touch surfaces included bed rail, bedside table, call button, and patients' bedside chart.

^dSkin sites included hands, chest, and groin.

**Fig. 2.** Cumulative percentage of methicillin-resistant *Staphylococcus aureus* (MRSA) contamination in rooms of 17 study participants and percentage of patients remaining the hospital.

evidence that interactions with personnel or equipment resulted in contamination of patient's skin or high-touch surfaces in rooms of hospitalized patients as all cultures collected immediately after observed interactions were negative. Rather, healthcare-associated pathogens, particularly MRSA, were initially recovered from the floor in rooms of newly admitted patients followed in a subset of patients by detection on sock bottoms, bedding, and high-touch surfaces. In 2 (66.7%) of 3 instances when MRSA was recovered from high-touch surfaces, an isolate with the same *spa* type had been recovered from the floor previously. These findings build upon a growing body of evidence suggesting that floors might

be an underappreciated source of pathogen dissemination not addressed by current control measures.^{2,9-13}

The fact that pathogens were recovered from the socks of patients suggests that one route of transfer may involve movement from socks to bedding or hands with subsequent transfer to high-touch surfaces by hands. The simulations with bacteriophage MS2 demonstrate the feasibility of this mechanism of transfer, either from socks directly to hands during sock removal or from socks to bedding and then to hands. Alternatively, we have previously demonstrated that fomites often contact hospital room floors and can serve as a vector for transfer to hands.¹⁰




Fig. 3. Concentration of bacteriophage MS2 recovered from different sites during 2 simulation protocols with or without slippers worn over socks and removed on sitting. Bacteriophage MS2 was inoculated onto the floor and allowed to air dry. A, participants stepped onto the contaminated area, sat on the bed, removed slippers (if present), lay down on the bed, touched the bedding adjacent to their feet, and touched the bedside table. B, participants stepped onto the contaminated area, sat on the bed, removed slippers (if present), removed socks, and touched the bedside table. PFU, plaque-forming unit.

Our results suggest that 2 simple interventions could be helpful in reducing the risk for acquisition of pathogens from hospital room floors. First, cleaning and disinfection of floors with a relatively dilute sodium hypochlorite disinfectant followed by operation of a UV-C room decontamination device after routine postdischarge cleaning resulted in consistent negative cultures of floors. In a previous study, cleaning and disinfection of floors with a quaternary ammonium disinfectant and/or operation of a UV-C room decontamination device were effective in reducing floor contamination.¹⁸ Second, in the simulation study, wearing slippers over hospital socks dramatically reduced transfer of bacteriophage MS2 from the floor to hands and to high-touch surfaces. Because extensive efforts to clean floors are not likely to be feasible in healthcare facilities, having patients wear slippers might be an easy measure to reduce the frequency of transfer of pathogens from the floor to hands and high-touch surfaces.

Our study has several limitations. We studied a relatively small number of patients from a single facility with excellent hand hygiene compliance rates. Additional studies are needed in other settings. Our results suggest but do not prove that shoes transferred healthcare-associated pathogens into rooms of patients. Previously, we demonstrated that shoes of personnel are frequently contaminated with pathogens, particularly MRSA.¹⁵ The demonstration of the effectiveness of slippers in reducing transfer from floors involved a simulation with healthy volunteers. Additional studies are needed with hospitalized patients who could have difficulty wearing slippers due to issues such as frailty or loss of visual acuity. We studied only 3 common bacterial pathogens and therefore our findings may underestimate the potential for importation of pathogens such as *Candida* spp and viruses, including severe acute respiratory syndrome coronavirus 2, that have been detected on floors.^{19–21} We did not closely observe interactions between patients and visitors who could also potentially serve as a source of importation of pathogens.²² Rectal cultures were not collected on admission to assess for VRE or *C. difficile* colonization. However, groin and skin swabs were collected. Previous studies have demonstrated that groin and skin cultures detect approximately half of patients colonized with VRE or *C. difficile*, including those most likely to shed organisms to the environment.^{23,24} Finally, we did not evaluate strategies that might reduce transfer of pathogens into patient rooms on shoes or on wheeled equipment. For example, it has been suggested that the use of a

UV-C device for shoe decontamination might reduce transfer of pathogens into rooms on shoes.²⁵ Improved cleaning and disinfection protocols for devices such as wheelchairs have also been suggested as a means to limit the transfer of pathogens by contaminated wheels.²⁶

In conclusion, our findings suggest that healthcare facility floors may be an underappreciated source of dissemination of healthcare-associated pathogens. Future studies are needed to obtain additional evidence regarding the role of floors in pathogen transmission and to evaluate potential control measures. Simple interventions such as improved floor cleaning and disinfection, having patients wear slippers, and cleaning of high-touch items that contact the floor deserve additional study.

Acknowledgments. We thank the nurses, environmental management services personnel, and patients at the Cleveland VA Medical Center who participated in the study.

Financial support. This work was supported by from the Department of Veterans Affairs (merit review grant no. CX001848) to C.J.D.

Conflicts of interest. C.J.D. has received research grants from Clorox, Pfizer, and PDI. All other authors report no conflicts of interest relevant to this article.

References

- Donskey CJ. Does improving surface cleaning and disinfection reduce healthcare-associated infections? *Am J Infect Control* 2013;41:S12–S19.
- Donskey CJ. Beyond high-touch surfaces: portable equipment and floors as potential sources of transmission of healthcare-associated pathogens. *Am J Infect Control* 2019;47S:A90–A95.
- Kanamori H, Rutala WA, Weber DJ. The role of patient care items as a fomite in healthcare-associated outbreaks and infection prevention. *Clin Infect Dis* 2017;65:1412–1419.
- Suwanarat N, Supple LA, Cadnum JL, Sankar T, Donskey CJ. Quantitative assessment of interactions between hospitalized patients and portable medical equipment and other fomites. *Am J Infect Control* 2017;45:1276–1278.
- John A, Alhmidhi H, Cadnum JL, Jencson AL, Donskey CJ. Contaminated portable equipment is a potential vector for dissemination of pathogens in the intensive care unit. *Infect Control Hosp Epidemiol* 2017;38:1247–1249.
- Jernigan JA, Siegman-Igra Y, Guerrant RC, Farr BM. A randomized cross-over study of disposable thermometers for prevention of *Clostridium difficile* and other nosocomial infections. *Infect Control Hosp Epidemiol* 1998;19:494–499.
- John AR, Alhmidhi H, Cadnum JL, Jencson AL, Gestrich S, Donskey CJ. Evaluation of the potential for electronic thermometers to contribute to

- spread of healthcare-associated pathogens. *Am J Infect Control* 2018; 46:708–710.
8. Eyre DW, Sheppard AE, Madder H, *et al.* A *Candida auris* outbreak and its control in an intensive care setting. *N Engl J Med* 2018;379:1322–1331.
 9. Koganti S, Alhmidi H, Tomas ME, Cadnum JL, Jencson A, Donskey CJ. Evaluation of hospital floors as a potential source of pathogen dissemination using a nonpathogenic virus as a surrogate marker. *Infect Control Hosp Epidemiol* 2016;37:1374–1377.
 10. Deshpande A, Cadnum JL, Fertelli D, *et al.* Are hospital floors an underappreciated reservoir for transmission of healthcare-associated pathogens? *Am J Infect Control* 2017;45:336–338.
 11. Rashid T, Vonville H, Hasan I, Garey KW. Mechanisms for floor surfaces or environmental ground contamination to cause human infection: a systematic review. *Epidemiol Infect* 2017;145:347–357.
 12. Galvin J, Almatroudi A, Vickery K, *et al.* Patient shoe covers: transferring bacteria from the floor onto surgical bedsheets. *Am J Infect Control* 2016; 44:1417–1419.
 13. Mahida N, Boswell T. Non-slip socks: a potential reservoir for transmitting multidrug resistant organisms in hospitals? *J Hosp Infect* 2016;94:273–275.
 14. Cadnum JL, Tomas ME, Sankar T, *et al.* Effect of variation in test methods on performance of ultraviolet-C radiation room decontamination. *Infect Control Hosp Epidemiol* 2016;37:555–560.
 15. Kanwar A, Thakur M, Wazzan M, *et al.* Clothing and shoes of personnel as potential vectors for transfer of healthcare-associated pathogens to the community. *Am J Infect Control* 2019;47:577–579.
 16. Alhmidi H, Cadnum JL, Koganti S, *et al.* Shedding of methicillin-resistant *Staphylococcus aureus* by colonized patients during procedures and patient care activities. *Infect Control Hosp Epidemiol* 2019;40:328–332.
 17. Tomas ME, Kundrapu S, Thota P, *et al.* Contamination of the skin and clothing of healthcare personnel during removal of personal protective equipment. *JAMA Intern Med* 2015;175:1904–1910.
 18. Mustapha A, Alhmidi H, Cadnum JL, Jencson AL, Donskey CJ. Efficacy of manual cleaning and an ultraviolet C room decontamination device in reducing healthcare-associated pathogens on hospital floors. *Am J Infect Control* 2018;46:584–586.
 19. Kumar J, Eilertson B, Cadnum JL, *et al.* Environmental contamination with *Candida* species in multiple hospitals including a tertiary-care hospital with a *Candida auris* outbreak. *Pathog Immun* 2019;4:260–270.
 20. Ye G, Lin H, Chen S, *et al.* Environmental contamination of SARS-CoV-2 in healthcare premises. *J Infect* 2020;81(2):e1–e5.
 21. Redmond SN, Dousa KM, Jones LD, *et al.* Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) RNA contamination of surfaces on a coronavirus disease 2019 ward and intensive care unit. *Infect Control Hosp Epidemiol* 2020. doi: [org/10.1017/ice.2020.416](https://doi.org/10.1017/ice.2020.416).
 22. Knighton SC, Zabarsky TF, Donskey CJ. Four moments for health-care facility visitor hand hygiene. *Am J Infect Control* 2020;48: 1412–1413.
 23. Sethi AK, Al-Nassir WN, Nerandzic MM, Donskey CJ. Skin and environmental contamination with vancomycin-resistant enterococci in patients receiving oral metronidazole or oral vancomycin treatment for *Clostridium difficile*-associated disease. *Infect Control Hosp Epidemiol* 2009;30:13–17.
 24. Donskey CJ, Sunkesula VCK, Jencson AL, *et al.* Utility of a commercial PCR assay and a clinical prediction rule for detection of toxigenic *Clostridium difficile* in asymptomatic carriers. *J Clin Microbiol* 2014;52: 315–318.
 25. Rashid T, Poblete K, Amadio J *et al.* Evaluation of a shoe sole UVC device to reduce pathogen colonization on floors, surfaces and patients. *J Hosp Infect* 2018;98:96–101.
 26. Jencson AL, Cadnum JL, Wilson BM, Donskey CJ. Spores on wheels: wheelchairs are a potential vector for dissemination of pathogens in healthcare facilities. *Am J Infect Control* 2019;47:459–461.

© The Author(s), 2021. Published by Cambridge University Press on behalf of The Society for Healthcare Epidemiology of America. This work is classified, for copyright purposes, as a work of the U.S. Government and is not subject to copyright protection within the United States.